ELSEVIER

Journal of Immunological Methods 205 (1997) 213–214

JOURNAL OF IMMUNOLOGICAL METHODS

Contents of volume 205 (1997)

Number 1 – 23 June 1997

Research reports

Human umbilical cord blood-derived eosinophils cultured in the presence of IL-3 and IL-5 respond to fMLP with $[Ca^{2+}]_i$ variation and O_2^- production
D.M. Zardini, P. Heuschling, A. Gallois, J.-L. Bueb and E.J. Tschirhart (Luxembourg) 1

Pilot scale purification of human monoclonal IgM (COU-1) for clinical trials
I. Tornøe, I.L. Titlestad, K. Kejling, K. Erb, H.J. Ditzel and J.C. Jensenius (Denmark) 11

A simple method for the propagation and purification of γδT cells from the peripheral blood of glioblastoma patients using solid-phase anti-CD3 antibody and soluble IL-2
T. Yamaguchi, Y. Fujimiya, Y. Suzuki, R. Katakura and T. Ebina (Japan) 19

A simple method for the preparation and purification of C1 complement cleaved β_2-microglobulin from human serum
M.H. Nissen, B. Johansen and O.J. Bjerrum (Denmark) 29

A specific and sensitive ELISA for measuring S-100b in cerebrospinal fluid
A.J.E. Green, G. Keir and E.J. Thompson (UK) 35

Importance of the linker in expression of single-chain Fv antibody fragments: optimisation of peptide sequence using phage display technology
D.J. Turner, M.A. Ritter and A.J.T. George (UK) 43

Flow cytometric quantitation of attachment and phagocytosis in phenotypically-defined subpopulations of cells using PKH26-labeled FcγR-specific probes
L. Pricop, J.E. Salmon, J.C. Edberg and A.J. Beavis (USA) 55

Correlation between the avidity of mouse-human chimeric IgG subclass monoclonal antibodies measured by solid-phase elution ELISA and biospecific interaction analysis (BIA)
N. McCloskey, M.W. Turner and D. Goldblatt (UK) 67

Large scale and clinical grade purification of syndecan-1 + malignant plasma cells
R.-X. Sun, Z.-Y. Lu, J. Wijdenes, J. Brochier, C. Hertog, J.-F. Rossi and B. Klein (France) 73

A sensitive enzyme immunoassay of human thyroid-stimulating hormone (TSH) using bispecific $F(ab')_2$ fragments recognizing polymerized alkaline phosphatase and TSH
K. Morimoto and K. Inouye (Japan) 81

Short communications

Multiple immunoblots after non-electrophoretic bidirectional transfer of a single SDS–PAGE gel with multiple antigens
B.T. Kurien and R.H. Scofield (USA) 91

The use of Apostain in identifying early apoptosis
C. Ferlini, A. Kunkl, G. Scambia and A. Fattorossi (Italy) 95

Book review 103

Corrigendum

Corrigendum to "Production of polyclonal antibody specific for human natriuretic peptide receptor B" [J. Immunol. Methods 194 (1996) 147–153]
K. Kitano, Y. Fukuda, K. Nagahira, T. Nasu, C. Noguchi, R. Izumi, K. Kawashima and T. Nakanishi (Japan) 105

Notes to authors 107

Number 2 – 14 July 1997

Research reports

Measurement of glucose consumption by hybridoma cells growing in hollow fiber cartridge bioreactors: use of blood glucose self-monitoring devices
R.C. Nayak and I.M. Herman (USA) 109

Organ culture of human lymphoid tissue. II. Marked differences in cytokine production and proliferation between slice and suspension cultures of human spleen
G. Skibinski, P. Hoffmann, A. Radbruch and K. James (UK, Germany) 115

Use of native and recombinant bactericidal/permeability-increasing proteins (BPI) as antigens for detection of BPI-ANCA
H. Schultz, E. Csernok, T.W. Johnston, C.M. Lockwood and W.L. Gross (Germany, UK) 127

A sensitive immunoassay to detect the α-chemokine GRO in rabbit blood and lung fluids
O. Kajikawa, M.C. Johnson II, R.B. Goodman, C.W. Frevert and T.R. Martin (USA) 135

Consistent transient transfection of DNA into non-transformed human and murine T-lymphocytes
R.Q. Cron, L.A. Schubert, D.B. Lewis and C.C.W. Hughes (USA) 145

A comparative study of optical techniques applied to particle-enhanced assays of C-reactive protein
J.L. Ortega-Vinuesa, J.A. Molina-Bolívar, J.M. Peula and R. Hidalgo-Alvarez (Spain) 151

A simple SDS–PAGE immunoblotting technique using an enhanced chemiluminescence detection system to identify polyclonal antibody responses to complex cryptosporidial antigen preparations following a monoclonal antibody retest and image overlay technique
S. Patel, J. McLauchlin and D.P. Casemore (UK) 157

Incorporation of cells into an ELISA system enhances antigen-driven lymphokine detection
J.T. Beech, T. Bainbridge and S.J. Thompson (UK) 163

Dual-label time-resolved immunofluorometric assay for simultaneous determination of pregnancy-associated plasma protein A and free β-subunit of human chorionic gonadotrophin
Q. Qin, M. Christiansen, T. Lövgren, B. Nørgaard-Pedersen and K. Pettersson (Finland, Denmark) 169

Production and characterization of monoclonal antibodies against the leukemia inhibitory factor low affinity receptor, gp190
V. Pitard, J.-L. Taupin, V. Miossec, F. Blanchard, M. Cransac, I. Jollet, A. Vernallis, K. Hudson, A. Godard, Y. Jacques and J.-F. Moreau (France, UK) 177

Detection of meningitis antigens in buffer and body fluids by ultrasound-enhanced particle agglutination
P. Jenkins, R.A. Barnes and W.T. Coakley (UK) 191

A novel, highly efficient peptide-HLA class I binding assay using unfolded heavy chain molecules: identification of HIV-1 derived peptides that bind to HLA-A* 0201 and HLA-A* 0301
T.L.R. Tan, A. Geluk, M. Toebes, T.H.M. Ottenhoff and J.W. Drijfhout (Netherlands) 201

Letter to the Editors

Cloning of cells with autologous feeder cells
A.W. Hügin (Switzerland) 211

Contents of volume 205 213

Notes to authors 215